



## **References for Neocate®**

### **2011**

1. Haas A. Development, Mealtime Dynamics and Nutrition: Use of an Amino Acid-Based Semi-Solid Supplement in a Child with Eosinophilic Esophagitis; Nutrition Practice Poster Exchange Clinical Nutrition Week 2011. *Nutr Clin Pract.* 2011;26(1):19(N36)

### **2010**

2. De Greef, et al. The influence of Neocate in Paediatric Short Bowel Syndrome on PN Weaning. *J Nutr Metab* 2010 pii: 297575. Epub 2010 May 31.

### **2009**

3. Becker JP. Cows Milk Allergy in a Preterm Infant; A Case Presentation. *ICAN: Infant Child & Adolescent Nutrition.* 2009;214-219.

### **2008**

4. Spergel JM, et al. 14 Years of Eosinophilic Esophagitis: Clinical Features and Prognosis. *J of Pediatr Nutr.* 2008;48:30-36.

### **2007**

5. Hill DJ, et al. The efficacy of amino acid based formula in relieving the symptoms of cows milk allergy-A systematic review. *Clin Exp Allergy.* 2007;(37):808-822.

### **2006**

6. Thomson M, et al. Effect of an amino acid-based milk-Neocate on gastro-oesophageal reflux in infants assessed by combined intraluminal impedance/pH, *Pediatr Allergy Immunol.* 2006; 19[4]:205-213.

### **2005**

7. Liacouras C, et al. Eosinophilic Esophagitis: A ten year experience in 381 children. *Clin Gastroenterol Hepatol.* 2005;3:1198-1206.
8. Faber MR et al, Allergic colitis presenting within the first hours of premature life, *Acta Paediatric* 2005;1514-1515.
9. Spergel JM, et al. Treatment of eosinophilic esophagitis with specific food elimination diet directed by a combination of skin prick and patch tests. *Ann Allerg Asthma Im.* 2005;95:336-343.

### **2004**

10. Arvola T, et al. Weaning to hypoallergenic formula improves gut barrier function in breast-fed infants with atopic eczema *J Pediatr Gastroenterol Nutr.* 2004;38(1): 92-96
11. Isolauri E et al. Use of an amino acid-based formula in the management of cow's milk allergy and multiple food protein intolerance in children. *J Allergy Clin Immunol.* 2004;113(2):s154 (518)
12. Hernandez-Trujillo VP, et al. Cow's milk allergy in a patient with hyper-IgE syndrome. *Ann Allerg Asthma Im.* 2004;92(4):469-474



**2003**

13. Hernandez-Trujillo VP, et al. Beneficial effect of an amino acid-based formula for neonatal Eosinophilic folliculitis associated with hypo-IgE Syndrome and cow's milk allergy. *J Allergy Clin Immunol.* 2003;111(2):s228 (638)
14. Heine R, et al. Effect of an amino acid based formula on gastroesophageal reflux in infants with persistent distress. *J Allergy Clin Immunol.* 2003;s102.A129
15. Markowitz JE, et al. Elemental diet is an effective treatment for eosinophilic esophagitis in children and adolescents. *Am J Gastroenterol.* 2003;98:777-782.

**2002**

16. De Boissieu D, et al. Allergy to extensively hydrolyzed cow's milk proteins in infants : Safety and duration of amino acid-based formula. *Pediatr.* 2002;141:271-3.
17. Serrano MS, et al. Nutrition support of infants with short bowel syndrome. *Nutrition.* 2002.18:966-970
18. Kanny G, et al. Use of an amino-acid-based formula in the treatment of cow's milk protein allergy and multiple food allergy Syndrome. *Allergie et immunologie.* 2002;34(3):82-4.
19. Dupont C, et al. Treatment of severe cow's milk protein allergy using Neocate. *Allergie et immunologie.* 2002;34(3):85-90.
20. Miele E, et al. Clinical response to Amino acid-based formula in neurologically impaired children with refractory Esophagitis. *J Pediatr Gastroenterol Nutr.* 2002;35:314-319
21. Amadi B. Role of food antigen elimination in treating children with persistent diarrhoea and malnutrition in Zambia. *J Paediatr Gastroenterol Nutr.* 2002 34: S54-56

**2001**

22. Andorsky DJ, et al. Nutritional and other postoperative management of neonates with short bowel syndrome correlates with clinical outcomes. *J Pediatr.* 2001;139:27-33
23. Niggemann B, et al. Prospective, controlled, multi-center study on the effect of an amino acid-based formula in infants with cow's milk allergy/intolerance and atopic dermatitis. *Pediatr Allergy Immunol.* 2001;12:78-82.
24. Moneret-Vautrin DA, et al. Milk protein hydrolysates: Hypoallergenic formulas and extensive hydrolysates: Allergologic basis for their use in the prevention and treatment of cow's milk allergy. *Archives de Pediatrie.* 2001;8(12):1348-57.

**2000**

25. Estep DC, et al. Treatment of infant colic with amino acid-based infant formula: a preliminary study. *Acta Paediatr.* 2000;89:22-7.
26. D'Netto MA, et al. Allergic gastroenteropathy in preterm infants. *J Pediatr.* 2000;137: 4: 480-486
27. Estep D C, et al. Colic in breast-milk-fed infants: treatment by temporary substitution of Neocate infant formula. *Acta Paediatr.* 2000;89:795-802



28. de Boissieu D, et al. Time course of allergy to extensively hydrolyzed cow's milk proteins in infants. *J Pediatr.* 2000;136:119-20.
29. Hill D, et al. Role of food protein intolerance in infants with persistent distress attributed to reflux esophagitis. *J Pediatr.* 2000;136:641-7.
30. Veereman-Wauters G, et al. Successful avoidance of total parenteral nutrition by administration of an amino acid based enteral formula (Neocate) in infancy. *Gastroenterology.* ,2000;31(2):s170.

**1999**

31. Hill D, et al. The natural history of intolerance to soy and extensively hydrolyzed formula in infants with multiple food protein intolerance. *J Pediatr.* 1999.135:118-21.
32. Isolauri E, et al. Breast-feeding of allergic infants. *J Pediatr.* 1999;134:27-32
33. Dupont C, et al. Early introduction of an amino acid-based vs protein hydrolysate formula in children with cow milk allergy: a randomised multicentre trial. *J Pediatr Gastroenterol Nutr.* 1999; 28:5: 589
34. Miele E, et al. Clinical response to amino acid-based enteral formula in neurologically impaired children with refractory Esophagitis. *J Pediatr Gastroenterol Nutr.*,1999;28(5):561.
35. Ammar F, et al. Allergy to protein hydrolysates; Report of 30 cases. *Archives de Pediatrie.* 1999;6:837-43

**1998**

36. Brewster D, at al. Short Bowel Syndrome, Intestinal Permeability and Glutamine. *J Pediatr Gastroenterol Nutr.* 1998; 27(5): 614-15.
37. Bines J, et al. Reducing parenteral requirement in children with short bowel syndrome: Impact of an amino acid-based complete infant formula. *J Pediatr Gastroenterol Nutr.* 1998;26:123-128

**1997**

38. Lake A.M. Beyond hydrolysates: Use of L-amino acid formula in resistant dietary protein-induced intestinal disease in Infants. *J Pediatr.*1997;131(5):658-60
39. Vanderhoof J.A et al. Intolerance to protein hydrolysate infant formulas: An under recognized cause of gastrointestinal symptoms in infants. *J Pediatr.* 1997;131:741-4.
40. de Boissieu D, et al. Allergy to extensively hydrolyzed cow milk proteins in infants: identification and treatment with an amino acid-based formula. *J Pediatr.* 1997;131:744-7.

**1996**

41. Isolauri E, et al. Combined skin prick and patch testing enhances identification of food allergy in infants with atopic dermatitis. *J Allergy Clin Immunol.* 1996;97:9-15
42. Kuzminskiene R. Nutritional management of multiple food protein intolerance induced severe atopic dermatitis in infants. *Allergy.* 1996;51(32):75(p94)



43. Murray RN, et al. Use of Neocate in infants with short bowel syndrome *20th Clinical Congress of the American Society for Parenteral and Enteral Nutrition. Washington DC 14-16, Jan 1996. Abstract 6*
44. Hill DJ, et al. Is multiple food intolerance the cause of reflux oesophagitis in distressed infants (abstract). *J Allergy Clin Immunol.*1996;(615) 336.
45. Murray ND, et al. Growth of infants who were treated with Neocate. *20<sup>th</sup> Clinical Congress Nutrition Practice Poster. Washington DC 14-16 1996. Abstract 5*

**1995**

46. Isolauri E, et al. Efficacy and safety of hydrolyzed cow milk and amino acid-derived formulas in infants with cow milk allergy. *J Pediatr.* 1995;127:550-7.
47. Hill DJ, et al. Challenge confirmation of late onset reactions to extensively hydrolyzed formulas in infants with multiple food protein intolerance. *J Allergy Clin Immunol.* 1995;96:386-94.
48. Kelly KJ, et al. Eosinophilic esophagitis attributed to gastroesophageal reflux: Improvement with an amino acid-based formula. *Gastroenterology.* 1995;109:1503-1512.

**1993**

49. Kelso JM, et al. Food protein-induced enterocolitis to casein hydrolysate formulas. *J Allergy Clin Immunol.* 1993;92:909-10

**1992**

50. Sampson HA, et al. Safety of an amino acid-derived infant formula in children allergic to cow milk. *Pediatrics.* 1992;90(3):463-465.

\*Most of the key studies have used Neocate Infant in a wide range of ages, from infancy through childhood. Many of the studies in children carried out before the development of Neocate One+ or Neocate Junior used Neocate because there were no AAF specifically designed to meet the nutritional requirements of children. It was as a consequence of studies such as these that the need for an age-specific feed for children was identified, and hence the development of Neocate One+ and Neocate Junior followed several years after the introduction of Neocate.



## Clinical References

### Nutricia North America Metabolic Products

#### Periflex Infant

1. Cleary MA, et al. Randomised controlled trial of essential fatty acid supplementation in phenylketonuria. *Eur J Clin Nutr.* 2006;60(7):915-20.
2. Agostoni C, et al. A randomized trial of long-chain polyunsaturated fatty acid supplementation in infants with phenylketonuria. *Dev Med Child Neurol.* 2006;48(3):207-12.
3. Greve LC, et al. Breast-Feeding in the Management of the Newborn with Phenylketonuria: A Practical Approach to Dietary Therapy. *J Am Diet Assoc.* 1994;94(3):305-9.
4. Mofidi S, et al. Growth And Plasma Phenylalanine And Tyrosine Concentrations Of Infants Treated With XPhe Analog Amino Acid Modified Infant Formula With Iron. Presented at the V International Congress of Inborn Errors of Metabolism, June 1-5, 1990.
5. Various authors. A Multicentre Study to Evaluate and Compare the Nutritional Adequacy of XP Analog and Lofenalac in the Dietary Management of Infants with Phenylketonuria (PKU). (1986-1989). Data on file.

#### Periflex Junior

1. Clear MA, et al. Randomised controlled trial of essential fatty acid supplementation in phenylketonuria. *Eur J Clin Nutr.* 2006;60(7):915-20.
2. Rose HJ, et al. Fat intakes of children with PKU on low phenylalanine diets. *J Hum Nutr Diet.* 2005;18(5):395-400.
3. Buist NRM, et al. A new amino acid mixture permits new approaches to the treatment of phenylketonuria. *Acta Paediatr Suppl.* 1994;407:75-7.

#### XPhe Maxamaid

1. MacDonald A, et al. Factors Affecting the Variation in Plasma Phenylalanine in Patients with Phenylketonuria on Diet. *Arch Dis Child.* 1996; 74: 412-417.
2. MacDonald A, et al. Feeding Problems in Young PKU Children. *Acta Paediatr.* 1994; 407: (Suppl) 73-74.
3. MacDonald A. Diet and Phenylketonuria: A Time for Change? *J Hum Nutr Diet.* 1994; 7:105-114.
4. Clinical evaluation report 1980-1982 by Prof. Frezal, Sick Children's Hospital Paris, available upon request.
5. Clinical evaluation report Dr. Sardhawalla, Royal Manchester Children's Hospital, available upon request.

#### XPhe Maxamum

1. Brown MC, et al Economic impact of feeding a phenylalanine-restricted diet to adults with previously untreated phenylketonuria. *J Intell Disabil Res.* 1999; 43:30-7.
2. Cleary MA, et al. Magnetic Resonance Imaging in Phenylketonuria: Reversal of Cerebral White Matter Change. *J Pediatr.* 1995; 127:251-255.
3. MacDonald A. Diet and Phenylketonuria: Time for Change? *J Hum Nutr Diet.* 1994; 7:105-114.
4. Wardley BL, et al. Clinical Trial of a Concentrated Amino Acid Formula for Older Patients with Phenylketouria (XP Maxamum). *Eur J Clin Nutr.* 1988; 42:81-86.
5. Taitz LS, et al. Zinc and Selenium Levels in Older Children with Phenylketonuria – Correction of Low Plasma Levels by a New Dietary Supplement Formulation (XP Maxamum). Data on file.



### XPhe Maxamum Drink

1. Shaw H, et al. Acceptability and Clinical use of a new Ready-To-Drink PKU product. Poster presented at annual meeting of GMDI (Genetic Metabolic Dietitians International), Atlanta, USA, 2006.

### Lanaflex

1. Schindeler S, et al. The effects of large neutral amino acids supplements in PKU: An MRS and neuropsychological study. *Mol Gen and Metab* 2007; 91:48-54.
2. Crowley C, et al. Clinical trial of “off diet” older phenylketonurics with a new phenylalanine-free product. *J Metal Deficiency Research* 1990; 34: 361-369.

### Phlexy-10

1. Moseley K, et al. Sapropterin Dihydrochloride (6R-BH4) and Maternal Phenylketonuria. *ICAN Infant, Child and Adolescent Nutrition* 2009; vol 1, no 5:262-6.
2. MacDonald A, et al. Protein substitutes for PKU: What's new? *J Inherit Metab Dis.* 2004; 27:363-371.
3. Rohr F, et al. Acceptability of a new modular protein substitute for the dietary treatment of phenylketonuria. *J Inherit Metab Dis.* 2001; 24:623-630.
4. Moseley K, et al. A new multi-module medical food product improves compliance in the treatment of phenylketonuria. Poster presented at annual meeting of SSIEM (Society for the Study of Inborn Errors of Metabolism), Cambridge, England, 2000.

### Add-Ins

1. Shaw H, et al. Phenylalanine (Phe) control in patients with phenylketonuria (PKU) consuming a novel metabolic medical food (Add-Ins™). Poster presentation at annual meeting of SSIEM (Society for the Study of Inborn Errors of Metabolism), Hamburg, Germany, 2007

### Monogen

1. Lessen R. Use of Skim Breast Milk for an Infant with Chylothorax. *ICAN* (2009); 1 (6):303-10
2. Ficicioglu C, et al. Very long-chain acyl-CoA dehydrogenase deficiency : The effects of acidnetal fat loading in a patient detected through newborn screening. *J Inherit Metab Dis.* 2009 Apr 4 (Epub ahead of print).
3. Izzard, ME et al. The use of Monogen on the conservative management of chylous fistula. *Otolaryng Head Neck.* 2007;136 (4 Suppl): S50-3.
4. Möller A, et al. Congenital intestinal lymphangiectasia: a rare differential diagnosis in hypoproteinemia in infants. *Klin Padiatrie.* 2006;218(4):224-5 (German).
5. Cormack, BE et al. Use of Monogen for pediatric postoperative chylothorax. *Ann Thorac Surg.* 2004; 77(1):301-5.
6. Ng PC, et al. Deceptive hyperbilirubinaemia in a newborn with familial lipoprotein lipase deficiency. *J Paediatr Child H.* 2001;37(3):314-6.



## Duocal

1. Illsinger S, et al. Branched chain amino acids as a parameter for catabolism in treated phenylketonuria. *Amino Acids*. 2005 Feb;28(1):45-50.
2. Plauth M, et al. Post-feeding hyperammonaemia in patients with transjugular intrahepatic portosystemic shunt and liver cirrhosis: role of small intestinal ammonia release and route of nutrient administration. *Gut*. 2000 Jun;46(6):849-55.
3. Wilson DC, et al. Randomised controlled trial of an aggressive nutritional regimen in sick very low birthweight infants. *Arch Dis Child Fetal Neonatal Ed*. 1997 Jul;77(1):F4-11.
4. Iacona G, et al. Case report: Multiple food allergy. *J Pediatr Gastroenterol Nutr* 1993; 16:472-473.
5. Morris R, et al. A comparison of the energy supplements Polycal and Duocal in cancer patients. *J Hum Nutr Diet*. 1990; 3: 171-176.

## **Clinical References and Articles for KetoCal, Ketogenic Diet and Modified Atkins Diet**

### **Studies Using KetoCal:**

#### **2011**

1. Kossoff EH, et al. Prospective study of the Modified Atkins Diet in combination with a ketogenic liquid supplement during the initial month. *J Child Neurol* 2011;26(2):147-51.

#### **2010**

2. Coppola G, et al. Ketogenic diet for the treatment of catastrophic epileptic encephalopathies in childhood. *Eur J Paediatr Neurol*. 2010 May;14(3):229-34. Epub 2009 Jul 25.
3. Wusthoff CJ, et al. The ketogenic diet in treatment of two adults with prolonged nonconvulsive status epilepticus. *Epilepsia*. 2010 Jun;51(6):1083-5. Epub 2009 Oct 20.

#### **2008**

4. Rubenstein JE. Use of the ketogenic diet in neonates and infants. *Epilepsia*. 2008 Nov;49 Suppl 8:30-2.